

# **Adaptive habitat selection by a single celled alga?**

A thesis presented to  
The Faculty of Graduate Studies  
of  
Lakehead University  
by

MARYJANE MOSES

In partial fulfillment of requirements for the degree of  
Master of Science in Biology

18/September/2013

© MaryJane Moses, 2013

## Summary

- 1) When populations have opportunity to occupy multiple habitats, individuals that move to habitats with higher mean fitness will expect to produce more descendants than will individuals that are incapable of adaptive habitat choice. Although adaptive movement is widely assumed, we lack understanding of how widely it might apply to motile but non-sentient organisms.
- 2) I used replicate populations of the single-celled alga, *Chlamydomonas reinhardtii*, to assess its ability to preferentially occupy habitat yielding the highest fitness. I pipetted different densities of *C. reinhardtii* into pairs of shaded and unshaded control Petri dishes filled with growth media. I estimated fitness of this ‘clonal’ species as per capita growth rate (the ratio of cell densities measured at time  $t + 1$  divided by the density at time  $t$ ). I used the estimates to predict the ideal-free distribution of cells expected in adjacent pairs of the two habitats.
- 3) I created pairs of adjacent shaded and unshaded habitats within two other sets of Petri dishes by covering one-half of each dish with black micromesh. One set of dishes contained unused media, the other set contained the same media in which the cells had been growing (used media). I pipetted algae into either half of these Petri dishes and let cells distribute between habitats for 12 h. I isolated the two halves of each dish and sampled the density of cells occupying each side. I compared the observed distribution with that predicted to test for an ideal-free distribution and calculated fitness to assess adaptive movement.
- 4) Fitness declined linearly with increasing density in both the light and shade controls, and was higher in light than in shade. When pipetted into the light side of dishes with unused

media, cells were more abundant in light than in shade, and there was no difference in fitness. But when pipetted into the shaded side in dishes with unused media, and in all treatments with used media, there was no significant difference in cell density between habitats even though fitness was usually higher in the light habitat.

- 5) It thus appears that the ability of *Chlamydomonas reinhardtii* to achieve an ideal free distribution, and more generally to move adaptively, is contingent not only on differences between habitats, but also on the mean quality of the environment in which habitat selection occurs. Regardless, the experiments demonstrate that a motile non-sentient species with simple sensory abilities is clearly capable of adaptive movement that enhances fitness.

## Lay Summary

Faculty and students in the Department of Biology are bound together by a common interest in explaining the diversity of life, the fit between form and function, and the distribution and abundance of organisms. The research reported here shows how habitat selection and adaptive movement influence population dynamics. I first demonstrate by theory why habitat selection should vary with density. I then describe experiments in which I measured per capita population growth rates of the single-celled alga, *Chlamydomonas reinhardtii*, in separate shaded and unshaded (light) habitats across a range of densities, and used them to predict the number of cells that one should observe in adjacent habitats. Fitness was higher in the light habitat than in the shade. Using these observations, theory predicts that all individuals should occupy the light habitat at low density, but as density increases individuals should increasingly occupy the shade. More generally, individual cells should move to habitats of higher mean fitness (adaptive movement). My experiments on *Chlamydomonas* demonstrated a rather novel form of adaptive movement that only partially confirmed the predictions. The fitness of cells released in the light habitat in rich environments was not different from that in shade because density was higher in light. Cells released in shade, and those released in poor environments, moved such that there were no differences in density between habitats, even though fitness was usually higher in light. Adaptive movement thus depends not only on the quality of the occupied habitat, but also on mean environmental quality.

## **Acknowledgments**

I am grateful for the continued support and guidance for this work from the curious and dedicated nature of my supervisor Dr. Douglas Morris. I thank M. Maki and S. Schroeder for their assistance with laboratory research, and A. Dupuch, W. Halliday, S. Vijayajan, and R. Buchkowski as members of the evolutionary ecology research team. I am also grateful to S. Hecnar, W. Qin, and B. Danielson for insightful advice and comments that helped enhance the value of this thesis. I thank Canada's Natural Sciences and Engineering Research Council (NSERC) for its continuing support of D. Morris' research program in evolutionary ecology. I acknowledge Lakehead University for additional scholarship support and collaborative efforts with Dr. W. Qin from the Biorefining Research Institute. Lastly, I thank my family and friends for encouraging me to pursue this research.

## Table of Contents

<b>Summary</b> .....	ii
<b>Lay summary</b> .....	iv
<b>Acknowledgments</b> .....	v
<b>Table of contents</b> .....	vi
<b>List of appendices</b> .....	vii
<b>List of tables</b> .....	viii
<b>List of figures</b> .....	ix
<b>Introduction</b> .....	1
<b>Material and methods</b> .....	2
ALGAL CULTURES.....	2
EXPERIMENTAL DESIGN.....	5
FITNESS & HABITAT SELECTION.....	6
PREDICTIONS & TESTS.....	7
DIFFUSION CONTROL.....	9
<b>Results</b> .....	10
UNUSED MEDIA.....	10
USED MEDIA.....	20
DIFFUSION TEST.....	29
<b>Discussion</b> .....	29
<b>References</b> .....	34

## List of Appendices

<b>Appendix 1: DENSITY-DEPENDENT HABITAT SELECTION THEORY.....</b>	<b>38</b>
<b>Appendix 2: HABITAT ISODAR THEORY.....</b>	<b>41</b>
<b>Appendix 3: BOLD’S BASAL MEDIUM.....</b>	<b>44</b>
<b>Appendix 4: QUANTIFYING CELL DENSITY.....</b>	<b>47</b>
<b>Appendix 5: LABORATORY &amp; EXPERIMENTAL PROTOCOLS.....</b>	<b>52</b>
<b>Appendix 6: PHOTOGRAPHS.....</b>	<b>55</b>
<b>Appendix 7: DIFFUSION.....</b>	<b>62</b>

## List of Tables

Table 1: Summaries of the relationships between per capita population growth rate (fitness) and cell density, of densities between the two sides of control dishes, and of densities between habitats in treatment dishes for unused media (geometric mean regression; 95% confidence intervals in parentheses). All regressions were statistically significant (bold lettering).....	15
Table 2: Comparisons of per capita population growth rates between the initial and alternate sides of Petri dishes containing controls and habitat-selection treatments in unused media. Bold lettering identifies statistically significant differences (non-IFD). Paired t-tests; two-tailed significance. ....	17
Table 3: Summaries of the relationships between per capita population growth rate (fitness) and cell density, of densities between the two sides of control dishes, and of densities between habitats in treatment dishes for used media (geometric mean regression; 95% confidence intervals in parentheses). Bold lettering identifies statistically significant differences.....	24
Table 4: Comparisons of per capita population growth rates between the initial and alternate sides of Petri dishes containing controls and habitat-selection treatments in used media. Bold lettering identifies statistically significant differences (non-IFD). Paired t-tests; two-tailed significance.....	26
Table A1: Stock solutions and volumes of each compound used in the recipe for modified Bold's basal medium. Original stock solution and refinements are listed as in Bold (1949) and Bell (1990).....	46
Table A2: Culturing and experimental protocol used to assess habitat selection by <i>Chlamydomonas reinhardtii</i> .....	54



Table A3: Comparisons of paired mean absorbancy values between the initial (dye added) and alternate (no dye) sides of Petri dishes at 12 hourly intervals. Degrees of freedom at 3 and 11 hours were reduced because I replaced one randomly chosen sample with a pure-media blank in order to calibrate the spectrophotometer for each of the two 96-well microplates evaluating absorbancy of the 192 samples (16 samples  $\times$  12 hours). Bold lettering identifies statistically significant differences. Paired t-tests; two-tailed significance.....64

## List of Figures

Figure 1: Digitally photographed <i>Chlamydomonas reinhardtii</i> cells [CC-2935, wildtype (-)] under 60X magnification using a phase-contrast setting on an inverted microscope (Olympus IX51, USA) in the Lakehead University Instrumentation Lab, July 2010.....	4
Figure 2: The relationships between per capita population growth rates (fitness) and <i>Chlamydomonas</i> cell density (millions of cells·ml <sup>-1</sup> ) in the control light (open circles) and control shade (filled circles) habitats with unused media ( $n = 30$ for each regression).....	13
Figure 3: The ‘isodar graphs’ of <i>C. reinhardtii</i> (millions of cells·ml <sup>-1</sup> ) living in unused media. a and b: habitat isodars comparing initiated and alternate sides from control dishes. c and d: regressions of density from the treatment dishes. Open data points represent experiments initiated in the light habitat, filled data points correspond to experiments initiated in the shade. Dashed lines represent the isodar predicted from comparisons between shade and light control dishes.....	19
Figure 4: The relationships between fitness (per capita population growth rates) and <i>Chlamydomonas</i> cell density (millions of cells·ml <sup>-1</sup> ) in the control light (open circles) and control shade (filled circles) habitats with used media ( $n = 26$ and $n = 29$ respectively).....	22
Figure 5: The ‘isodar graphs’ of <i>C. reinhardtii</i> (millions of cells·ml <sup>-1</sup> ) living in used media. a and b: habitat isodars comparing initial and alternate sides of control dishes. c and d: regressions of density from the treatment dishes. Open data points represent experiments initiated in the light habitat, filled data points correspond to experiments initiated in the shade. Dashed lines represent the isodar predicted from comparisons between shade and light control dishes.....	26

Figure A1: An illustration of ideal-free habitat selection with logistic population growth. Ideal-free habitat selectors achieve equal expectations of fitness ( $W$ ) in habitats 1 and 2 (dashed horizontal lines), but at different population densities ( $N_1$ and $N_2$ , dotted horizontal lines).....	40
Figure A2: An example of an ideal-free habitat isodar (the distribution of individuals between two habitats such that mean fitness is equal in each) that emerges when fitness declines linearly with increasing density ( $N$ ) in habitats 1 and 2 as in figure A1. The dashed lines correspond to the carrying capacities in each habitat.....	43
Figure A3: Regressions used to calibrate cell densities (millions of cells·ml <sup>-1</sup> ) from optical densities (absorbancy at 665 nm) at the start of the light cycle for both light (a: open squares) and shaded (b: filled squares) habitats ( $n = 18$ each). OD = optical density.....	49
Figure A4: Regressions used to calibrate cell densities (millions of cells·ml <sup>-1</sup> ) from optical densities (absorbancy at 665 nm) at the end of the light cycle (12 h) for both light (a: open squares) and shaded (b: filled squares) habitats ( $n = 18$ each). OD = optical density.....	51
Figure A5: Photograph illustrating an example of the random placement of control (fully covered by shade and fully exposed to light) and habitat-selection Petri dishes (half shaded) used to assess habitat selection by <i>Chlamydomonas reinhardtii</i> . Each of the three replicate dishes contains a different algal density. Image taken at 12 h.....	57
Figure A6: Photograph of a Petri dish illustrating habitat selection by <i>Chlamydomonas reinhardtii</i> cells released in the light habitat (right-hand side) with opportunity to occupy shade (unused media). Initial density at time zero = 0.85 million cells·ml <sup>-1</sup> . Image taken at 12 h.....	59
Figure A7: Photograph of immobilized <i>Chlamydomonas reinhardtii</i> cells in a haemocytometer chamber used to calibrate optical densities. Living cells are dark green. Dead cells (not counted) appear without complete cellular structure or vibrancy (Leboffe & Pierce 2005).....	61

## 1    **Introduction**

2    The vast majority of organisms compete for resources in spatially and temporally stochastic  
3    environments where density-dependent habitat selection can modulate population dynamics,  
4    species-interactions, and community structure (Rosenzweig 1981; Morris 2011). When  
5    organisms live in heterogeneous environments, those individuals which remain in or move to  
6    areas of high fitness will expect to produce more descendants than will individuals lacking such  
7    adaptive potential (Holt 1985; Abrams 2000). Classical theory imagines that the movement of  
8    individuals among habitats that differ in suitability should equalize mean fitness (an ideal free  
9    distribution [IFD]; Fretwell & Lucas 1969). The IFD assumes, however, that organisms possess  
10   perfect information and move only to increase fitness (Milinski & Parker 1991; Hugie & Grand  
11   1998). Somewhat less ‘perfect’ organisms are nevertheless likely to move to, or remain in, areas  
12   of higher fitness (adaptive movement: Abrams, Cressman & Křivan 2010) but not necessarily  
13   equalize mean fitness amongst habitats (Cressman & Křivan 2012).

14        Experiments that match variation in habitat quality with the sensory capabilities and  
15   motility of organisms should be able to detect adaptive movement. Such experiments will be  
16   most effective if they can be replicated under strictly controlled conditions and if they can  
17   independently assess fitness and density. Thus, I ask whether *Chlamydomonas reinhardtii* (Fig.  
18   1), a motile single-celled alga with phototactic and chemotactic abilities (Harris et al. 2009), can  
19   achieve adaptive movement.

20        I begin by describing how I manipulated population densities of *C. reinhardtii* in order to  
21   obtain replicated estimates of fitness (per capita population growth rates) in shaded and unshaded  
22   (light) habitats. I use the relationships between fitness and density to predict the expected

distributions of cells in the two habitats assuming ideal-free habitat selection. I then describe experiments where I covered one-half of a set of Petri dishes with micromesh to create adjacent pairs of shaded and unshaded habitats, and assess whether the algae preferentially occupied the habitat yielding the highest fitness. I contrast the observed patterns of distribution and fitness with those predicted from theory and conclude by discussing the evidence supporting adaptive movement, and how it might be constrained in environments of low mean quality.

## **Materials and methods**

### **ALGAL CULTURES**

Pure batch cultures of wild-type bi-flagellate *Chlamydomonas reinhardtii* (strain CC-2935, *Chlamydomonas* Center, Duke University, NC) were grown anaerobically in Erlenmeyer flasks containing modified Bold's basal medium (Appendix 3; Bold, 1940; Bell, 1990). Cultures were gently aerated with sterile filtered air at ambient CO<sub>2</sub> levels (AIRPUMP 702A, Rena®) using 5-ml glass Pasteur pipettes and flexible plastic tubing (C-Flex tubing, No. 06422-07, Cole Parmer). All cultures were grown synchronously in a growth chamber (ThermoScientific, Model no. 845, CA, USA) set on a 12 h light-dark cycle maintained at 23°C (±1.0°C). Mean (and standard deviation) light intensity obtained from measurements (Amprobe LM631A, WA, USA) taken every five minutes at 12 positions in the growth chamber over one hour (repeated four times at 3-h intervals,  $n = 144$ ,) was 2120 (±160) *lux*.

Cultures were started from single colonies grown on routinely transferred agar plates as described by Harris et al., (2009). Starting cultures were grown in 75 ml of fresh media in 250-ml Erlenmeyer flasks until they reached mid-log-phase density (1-5 million cells·ml<sup>-1</sup>) after five to seven days (Appendix 5; Table A2, p. 54). Aliquots (25 ml) were transferred into 300 ml of

46

47

48

49

50

51

52

53 Figure 1: Digitally photographed *Chlamydomonas reinhardtii* cells [CC-2935, wildtype (-)]  
54 under 60X magnification using a phase-contrast setting on an inverted microscope (Olympus  
55 IX51, USA) in the Lakehead University Instrumentation Lab, July 2010.

56

57

58

59

60

61

62


63

64

65

66

67



68

69

70

71

72

media in one litre Erlenmeyer flasks for an additional growth period (7-10 d). Culture densities were adjusted by dilution to achieve randomly allocated target densities, then centrifuged (Sorvall RC 6 Plus, No. 46910, Thermo Scientific, USA) to pelletize the cells (Harris et al. 2009). Cell pellets were washed and re-suspended in 100 ml of fresh media in 500-ml Erlenmeyer flasks, then acclimated for a further 36 h before experimental use (as recommended by Harris et al. 2009).

Optical densities were estimated from spectrophotometer absorbancy readings at 665 *nm* on a microplate spectrophotometer (xMark™ Microplate Absorbance Spectrophotometer #168-1150, Bio-Rad, USA) calibrated with one blank well containing pure media on each microplate. Optical densities of diluted cell cultures were converted to cell densities (millions of cells·ml<sup>-1</sup>) with calibration curves from haemocytometer cell counts of immobilized samples (Appendix 4, pp. 47-51; Fig. A7, p. 61).

## EXPERIMENTAL DESIGN

Experiments were conducted in glass Petri dishes (100 x 15 mm, No. 89000-304, VWR® International). All Petri dishes were placed inside 10-cm tall black cardstock cylinders to ensure that cultures received uniform light intensity from the bank of growth lights overhead. Removable partitions made of wax-based modelling clay (NDC57, Polyform Products Co. Illinois, USA) were placed across the middle of each Petri dish to create two equal-sized halves (habitats). Habitats designated as ‘shaded’ were overlaid with double-layered black fibreglass micromesh (mesh size 0.25 mm<sup>2</sup>, FCS7350-A, Saint Gobain, CA). The mesh on top of the Petri dish lids reduced the mean (and standard deviation) light intensity from 2120 (±160) to 240



( $\pm 35$ ) *lux* ( $n = 144$ ). Unshaded Petri dishes (light habitats) were fully exposed to the mean light intensity of 2120 ( $\pm 160$ ) *lux*.

Fifteen ml of *C. reinhardtii* culture was pipetted into one-half (the initial half) of the partitioned Petri dishes (15 ml of unused or used media without algae was pipetted into the alternate half). Partitions were removed and cells were free to move throughout the Petri dish for one full photoperiod (12 h). Two controls and two treatments, consisting of three different densities (12 Petri dishes), were tested simultaneously (Figure A5, p. 57). ‘Light’ controls were unshaded; shade controls were fully covered with mesh. One-half of each dish was covered by mesh in the two habitat-selection treatments. One treatment consisted of cell-culture pipetted into light halves of the three replicate dishes, the other consisted of cell-culture pipetted into shaded halves. All cell-culture transfers, positions in the growth chamber, and the order of processing samples were determined randomly in advance using a random number generator (R Core Development Team 2008).

## FITNESS AND HABITAT SELECTION

Cell densities in each habitat were quantified at time zero, and at 12, 24, and 48 hours after the start of each experiment. At the end of the first photoperiod (12 h), after cells had opportunity to select between habitats, new partitions were inserted between the habitats and remained in place for the duration of the experiment (Appendix 5; Fig. A6, p. 59). Each half of every Petri dish was gently aerated at each sampling interval to ensure accurate and repeatable density estimates within habitats. Eight-220  $\mu$ l samples from each half were drawn with an eight-tip micropipette oriented parallel to, and equidistant from (225 mm), the partition. All samples were then

117 pipetted into 96-well microplates (DL-3571172, BD Falcon, CA, USA) and the mean of three  
118 optical densities (absorbancy) recorded at 665 *nm* on the spectrophotometer. One ‘blank’ well  
119 containing pure media was used to calibrate the readings obtained from each microplate.  
120 Densities recorded at 12 h ( $N_{12}$ ) were used to determine the distribution achieved by habitat  
121 selection. Cells divide only during the dark part of the cycle (Harris et al. 2009), so I estimated  
122 fitness as the per capita population growth rate achieved between 24 and 48 h (density at 48 h  
123 [ $N_{48}$ ] divided by the density at 24 h [ $N_{24}$ ]). Use of the 24-48 h time period guarantees that the  
124 growth estimates represent fitness achieved after the cells had completed any habitat choice  
125 (division between h 12 and 24 would include fitness obtained by cells that occupied both habitats  
126 during the previous photosynthesis and habitat-selection phase of the experiment).

127 I conducted two sequential sets of experiments using two different kinds of media in  
128 order to assess the role of mean environmental quality on habitat selection. I used freshly  
129 prepared (unused) media with a full suite of nutrients in the first set of experiments. In the  
130 second set, I tested used media (media remaining after I removed cells by centrifugation) from  
131 cultures grown at equivalent densities (and for equal durations) as in the experimental cultures. I  
132 tested 15 experimental densities in each set of experiments (and for both the shaded and  
133 unshaded controls, as well as the two treatments; 120 Petri dishes yielding 240 estimates of  
134 density and fitness).

## 136 PREDICTIONS AND TESTS

137 I designed my experiments to assess whether motile organisms with sensory capability can use  
138 those abilities to remain in, or move to, habitats with higher mean fitness (adaptive movement).

I reasoned that an absence of covarying densities between the habitats would rule out density-dependent habitat selection. If the density in one habitat depended on that in the other, if the pattern of density mirrored that predicted from the control habitats, and if there was no difference in mean fitness between them, then I can conclude that *Chlamydomonas* habitat selection achieved an IFD. But if density in one habitat depended on that in the other, if mean fitness was higher in the rich (light) habitat, and if cells preferentially moved to or remained in the light, then I can conclude that *Chlamydomonas* is capable of adaptive habitat selection, but incapable of achieving an IFD.

I estimated the relationships between per capita population growth rates and cell density by geometric mean regressions in R with the lmodel2 package (2.15.2; R Development Core Team 2008). I used those functions (from the controls only) to predict the densities in each habitat such that mean fitness would be the same in each (the habitat isodars, Morris 1987; 1988; an IFD; Appendices 1 & 2, pp. 38-43). I then contrasted the actual relationships from the habitat-selection treatments with that predicted from the controls. I evaluated the fit of the models to the data by verifying that the distributions of residuals were not different from that expected assuming a normal distribution. I concluded the experiments by using paired t-tests to assess whether fitness was different between the initial and alternate halves of the Petri dishes in each experiment. Significance tests for those comparisons were two-tailed because the IFD predicts equal fitnesses and is silent as to which side of the dishes should attain higher fitness by a non-habitat selecting species.

## 161 DIFFUSION CONTROL

162 My tests for adaptive habitat selection assume that the light versus shade treatments created rich  
163 and poor habitats respectively, and that algal distributions differed from that expected by  
164 diffusion of the media between the two halves of the Petri dishes. If the media diffuses more  
165 slowly than the time-scale of habitat selection, then habitat-selection for light versus shade might  
166 be confounded by differences in nutrient concentrations.

167 I tested for this possibility by pipetting a 15-ml solution of media coloured with a single  
168 concentration of non-reactive dye (10 $\mu$ L of dye in 50 ml of media, Bio-Safe coomassie blue  
169 stain, #101-0786, BioRad, USA) into one half of 12 different Petri dishes. I pipetted 15 ml of  
170 standard media into the other half of the dishes. I evaluated the time-dependent pattern of  
171 diffusion by extracting eight-220  $\mu$ l paired samples from each side of 12 replicated Petri dishes  
172 at one-hour intervals for 12 hours (eight samples from each side of one dish at each interval, each  
173 dish used once only). I pipetted the samples into a 96-well microplate and recorded three  
174 absorbancy values at 665 *nm* on the spectrophotometer (this wavelength is close to the coomassie  
175 blue absorption maximum of 595 $nm$ , Syrový & Hodný 1991). I calculated the mean of the three  
176 values and assessed whether there were differences between the two halves of the dish with  
177 paired-t tests ( $n = 8$  for each of the 12 tests; Table A3, p. 64). I reasoned that the time when I  
178 was unable to detect a difference in absorbancy would correspond with that required for  
179 diffusion to ‘homogenize’ the media.

180

181

## Results

### UNUSED MEDIA

Per capita population growth rate was higher in the control light than in the control shade. Growth rates declined with population density linearly, and in parallel, in both habitats (equations A and B in Table 1, Fig. 2).

In order to assess whether or not the habitat-selection experiments yielded an IFD, I set equations (A) and (B) equal to one another

$$2.37 - 1.20 \times (\text{Density in light}) = 1.87 - 1.29 \times (\text{Density in shade}) \quad (1)$$

and solved for the habitat isodar as the density of cells occupying the light habitat,

$$\text{Density in light} = 0.42 + 1.08 \times \text{Density in shade} \quad (2)$$

(dashed line, Fig. 3c, d). According to the isodar, if cells select between the two habitats according to an ideal free distribution, then they should occupy only the light habitat at densities below 0.42 million cells·ml<sup>-1</sup>, then become ever more evenly distributed between habitats with increasing population size.

All regressions comparing cell densities between sides and habitats in unused media were statistically significant (Equations C-F; Table 1). Intercepts were not different from zero and slopes were not different from one in all comparisons of density between sides of control dishes (Table 1). Similarly, there was no difference in the mean per capita population growth rates between initial and alternate sides of the control Petri dishes (Table 2). Cells given a choice between the two identical halves of the control dishes followed an ideal free distribution.

When released in light habitat, there was no difference in mean fitness of *Chlamydomonas* between light and shade habitats (Table 2), but the resulting isodar departed from that predicted from the control dishes (intercept not different from zero, slope larger than unity (Fig 3c)). Cells did not achieve an ideal free distribution when released in the shaded half of the habitat-selection dishes. There was no difference in density between habitats (Table 1, Fig 3d) even though fitness was higher in the light habitat (Table 2).

221

222

223

224

225

226

227

228 Figure 2: The relationships between per capita population growth rates (fitness) and  
229 *Chlamydomonas* cell density (millions of cells·ml<sup>-1</sup>) in the control light (open circles) and  
230 control shade (filled circles) habitats with unused media ( $n = 30$  for each regression).

231

232

233

234

235

236


237

238

239

240

241



242

243

244

245



246

247

248

249

250

251

252

253 Table 1: Summaries of the relationships between per capita population growth rate (fitness) and  
254 cell density, of densities between the two sides of control dishes, and of densities between  
255 habitats in treatment dishes for unused media (geometric mean regression; 95% confidence  
256 intervals in parentheses). All regressions were statistically significant (bold lettering).

257

258

259

260

261

262

263

264

265

266

267

268

Comparison	Regression equation	<i>F</i>	df	<i>P</i>	Eq
<b>Per capita population growth rate and density (controls)</b>	$W_{light} = 2.37 (\pm 0.2) - 1.20 (\pm 0.3) \times N_{light}$	44.0	1, 28	<b>&lt;0.001</b>	A
	$W_{shade} = 1.87 (\pm 0.2) - 1.29 (\pm 0.4) \times N_{shade}$	6.4	1, 28	<b>0.02</b>	B
<b>Density between sides (controls)</b>	$N_{Alternate\ light} = -0.10 (\pm 0.3) + 0.82 (\pm 0.4) \times N_{Initial\ light}$	9.3	1, 13	<b>0.01</b>	C
	$N_{Alternate\ shade} = -0.02 (\pm 0.2) + 0.96 (\pm 0.4) \times N_{Initial\ shade}$	8.7	1, 13	<b>0.01</b>	D
<b>Density between habitats</b>	$N_{Alternate\ shade} = -0.03 (\pm 0.1) + 0.54 (\pm 0.2) \times N_{Initial\ light}$	21.5	1, 13	<b>&lt;0.001</b>	E
	$N_{Alternate\ light} = -0.03 (\pm 0.2) + 1.05 (\pm 0.4) \times N_{Initial\ shade}$	26.6	1, 13	<b>&lt;0.001</b>	F

269

270

271

272

273

274

275

276

277

278

279

280

281

282 Table 2: Comparisons of per capita population growth rates between the initial and alternate  
283 sides of Petri dishes containing controls and habitat-selection treatments in unused media. Bold  
284 lettering identifies statistically significant differences (non-IFD). Paired t-tests; two-tailed  
285 significance.

286

287

288

289

290

291

292

293

294

295

296

297

Side	Habitat	Mean fitness	Paired $T$	$P$
Initial	Light	1.56	1.92	0.08
Alternate	Light	1.77		
Initial	Shade	1.37	-1.32	0.21
Alternate	Shade	1.19		
Initial	Light	1.55	-1.8	0.09
Alternate	Shade	1.35		
Initial	Shade	1.10	3.97	<b>0.001</b>
Alternate	Light	1.79		

298

299

300

301

302

303

304

305

306

307

308

309

310

311 Figure 3: The ‘isodar graphs’ of *C. reinhardtii* (millions of cells·ml<sup>-1</sup>) living in unused media.  
312 a and b: habitat isodars comparing initial and alternate sides from control dishes. c and d:  
313 regressions of density from the treatment dishes. Open data points represent experiments  
314 initiated in the light habitat, filled data points correspond to experiments initiated in the shade.  
315 Dashed lines represent the isodar predicted from comparisons between shade and light control  
316 dishes.

317

318

319

320


321

322

323

324

325



326

327

328

## 329 USED MEDIA

330 Again, as in unused media, per capita population growth rate was higher in the control light than  
 331 in the control shade. Growth rates in controls declined with population density linearly, but  
 332 much more slowly in shade than in light (equations G and H in Table 3, Fig. 4). The maximum  
 333 per capita population growth in the control light was slightly lower than in unused media, as was  
 334 the decline with density (Table 3; contrast Fig. 2 with Fig. 4). There was no difference in fitness  
 335 between paired halves of control dishes in light (Table 4), but the regression comparing paired  
 336 densities in the two sides of the dishes was not significant (Fig. 5a, no evidence for density-  
 337 dependent habitat selection, and thus, not an IFD). Fitness and density in the initial shade side  
 338 exceeded that in the alternate shaded side (Table 4, Fig. 5b). Contrary to unused media, cells in  
 339 used-media controls failed to achieve an IFD. Nevertheless, I set equations (G) and (H) equal to  
 340 one another,

$$341 \quad 2.15 - 0.82 \times (\text{Density in light}) = 1.27 - 0.39 \times (\text{Density in shade}) \quad (3)$$

342 and solved the expected isodar as the cell density in the light habitat,

$$343 \quad \text{Density in light} = 1.07 + 0.48 \times \text{Density in shade} \quad (4)$$

344 (dashed line, Fig 5c, d). According to equation (4), if cells select between the two habitats  
 345 according to an ideal free distribution, then they should occupy only the light habitat at densities  
 346 below 1.07 million cells·ml<sup>-1</sup>. Then, for each increase in population size, approximately half as  
 347 many cells should occupy the lighted half of the dish as occupy the shaded half.

348 Neither habitat-selection treatment with used media produced patterns in habitat densities  
 349 different from linear one-to-one relationships with zero intercepts and slopes of unity (Table 3,

350

351

352

353

354

355

356

357 Figure 4: The relationships between fitness (per capita population growth rates) and  
358 *Chlamydomonas* cell density (millions of cells·ml<sup>-1</sup>) in the control light (open circles) and  
359 control shade (filled circles) habitats with used media ( $n = 26$  and  $n = 29$  respectively).

360

361

362

363

364

365

366


367



368

369

370



371

372

373

374

375

376

377

378

379

380

381

382 Table 3: Summaries of the relationships between per capita population growth rate (fitness) and  
383 cell density, of densities between the two sides of control dishes, and of densities between  
384 habitats in treatment dishes for used media (geometric mean regression; 95% confidence  
385 intervals in parentheses). Bold lettering identifies statistically significant differences.

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

Comparison	Regression equation	<i>F</i>	df	<i>P</i>	Eq
Per capita population growth rate and density (controls)	$W_{light} = 2.15 (\pm 0.2) - 0.82 (\pm 0.2) \times N_{light}$	32.6	1, 25	<b>&lt;0.001</b>	G
	$W_{shade} = 1.27 (\pm 0.1) - 0.39 (\pm 0.1) \times N_{shade}$	13.2	1, 27	<b>0.001</b>	H
Density between sides (controls)	$N_{Alternate\ light} = 0.12 (\pm 0.3) + 0.67 (\pm 0.4) \times N_{Initial\ light}$	3.5	1, 13	0.08	I
	$N_{Alternate\ shade} = -0.06 (\pm 0.2) + 1.32 (\pm 0.4) \times N_{Initial\ shade}$	9.6	1, 13	<b>&lt;0.01</b>	J
Density between habitats	$N_{Alternate\ shade} = -0.12 (\pm 0.2) + 0.91 (\pm 0.4) \times N_{Initial\ light}$	14.2	1, 13	<b>&lt;0.01</b>	K
	$N_{Alternate\ light} = -0.15 (\pm 0.2) + 0.84 (\pm 0.4) \times N_{Initial\ shade}$	7.4	1, 13	<b>0.02</b>	L

404

405

406

407

408

409

410

411 Table 4: Comparisons of per capita population growth rates between the initial and alternate  
412 sides of Petri dishes containing controls and habitat-selection treatments in used media. Bold  
413 lettering identifies statistically significant differences (non-IFD). Paired t-tests; two-tailed  
414 significance.

415

416

417

418

419

420

421

422

423

424

425

426

Side	Habitat	Mean fitness	Paired <i>T</i>	<i>P</i>
Initial	Light	1.56	1.82	0.09
Alternate	Light	1.91		
Initial	Shade	1.12	-2.20	<b>0.05</b>
Alternate	Shade	1.04		
Initial	Light	1.81	-5.01	<b>&lt;0.001</b>
Alternate	Shade	1.04		
Initial	Shade	1.12	3.97	<b>&lt;0.001</b>
Alternate	Light	2.16		

427

428

429

430

431

432

433

434

435

436

437

438

439

440 Figure 5: The ‘isodar graphs’ of *C. reinhardtii* (millions of cells·ml<sup>-1</sup>) living in used media.

441 a and b: habitat isodars comparing initial and alternate sides of control dishes. c and d:

442 regressions of density from the treatment dishes. Open data points represent experiments

443 initiated in the light habitat, filled data points correspond to experiments initiated in the shade.

444 Dashed lines represent the isodar predicted from comparisons between shade and light control

445 dishes.

446

447


448

449

450

451

452



454

455

456

457 Fig. 5c & d). Per capita population growth rates were significantly greater in light regardless of  
 458 which habitat the cells were released into (Table 4). Although there was no clear evidence of

differences in density between habitats, the pattern in the residuals suggests a possible preference for the light habitat at low density (Fig. 5c & d). This preference, if real, is consistent with the higher per capita population growth rates observed in that habitat at low density (Fig. 4).

## DIFFUSION

Mean absorbancy by dye was dramatically greater on the initial side of the dishes until about hour 4, after which absorbancy was more-or-less homogeneous in both halves of the dishes (Appendix 7, p. 62). It is thus reasonable to assume, for much of the 12-h time course available for habitat selection, 1, that media did not differ between light and shaded habitats, and 2, that differences between habitats were thus caused mainly by differences in light intensity.

## Discussion

Habitat selection operates through the process of adaptive movement such that individuals can increase fitness by moving to a different habitat. Motile organisms should thus evolve sensory capabilities that enable preferential occupation of habitats yielding higher fitness than others. The experiments reported here document that even simple organisms are capable of adaptive movement that yield repeated, but often less than ideal, patterns of habitat selection.

Adaptive habitat selection was best demonstrated in the experiments with unused media. When cells were released in either light or shade controls, there was no difference in density or fitness between the two halves of the dishes. But when cells were released in the light habitat and given the opportunity to move to shade, density was greater in the light habitat even though



my diffusion test showed that nutrient concentrations should have equilibrated between habitats. Cells released in light thus attained an ideal free distribution (fitness was not different between habitats), but the densities were not those predicted from the controls.

Conversely, there was no detectable difference in density between habitats when cells were exposed to used media, or released in the shade with unused media, even though mean fitness was higher in the light than in the shade in all treatments where cells demonstrated density-dependent habitat selection. How can adaptive movement account for these differing results?

Although the experiments maintained constant differences between light and shade habitats in light intensity, cell division was likely limited by local resource depletion as cells metabolized nutrients (Tilman 1990). Thus, diffusion of nutrients from used versus unused media should influence per capita population growth. In rich environments where unused media diffuses between habitats, cells in the light habitat are likely to receive adequate nutrients to maintain high photosynthetic output and reproduction. In poor (used media) environments, however, the availability of nutrients may limit photosynthetic efficiency and cell division such that there is little advantage in choosing one habitat over the other.

Regardless, fitness at most densities was higher in light habitat than in shade, and one should expect, *ceteris paribus*, a preference for light by phototactic algae. All is not quite equal, however, because cells living at high light intensities are exposed to increased oxidation of photosystems that require expensive cellular repairs (Harris et al. 2009). It is thus possible that the costs of oxidative stress in light limit the otherwise density-dependent benefits associated with occupying that habitat. Once restricted to the light habitat at the end of the 12-h selection phase, the cells might then invest in physiological adjustments to increase protection and

photosynthetic efficiency (Poulin et al. 2009). Those adjustments could then materialize as higher mean fitness (between 24 and 48 h) than the cells initially ‘anticipated’ during oxidative stress.

Some readers might question whether the chemotactic ability of *C. reinhardtii* also accounts for the apparent inability to preferentially occupy the habitat (light) yielding high fitness. I suspect not because diffusion should have equalized nutrient concentrations between habitats long before cells completed habitat selection. It is thus difficult to imagine chemotactic cues that, in my experiments, would consistently ‘attract’ cells to one habitat or the other.

When cells are released in light, the difference in movement responses between used and unused media are consistent with Křivan et al.’s (2008) prediction that the probability of emigration decreases with the increased suitability of the initial habitat. Cells appear to choose one habitat over another only when there is a strong signal of differences in quality between them that can overcome any costs associated with habitat selection. A similar pattern of apparently adaptive ‘non-movement’ occurs in laboratory populations of rotifers (*Brachionus calyciflorus*) which decrease speed and increase turning frequencies when occupying a high quality habitat (Kuefler, Avgar & Fryxell 2013). *C. reinhardtii* appears to possess a movement strategy similar to that of the rotifers. Cells living under diminished light switch stochastically between synchronous and asynchronous flagellar beating patterns that produce intervals of straight swimming with abrupt re-orientations (Poulin et al. 2009). Such simple decision making can likely pay substantial dividends in fitness because the swimming pattern increases the probability of contacting higher-quality habitat. The caveat is that *Chlamydomonas* may only be able to reap those dividends when resources are sufficiently abundant to maximize their photosynthetic capacity.

Algal habitat selection is further complicated by an ability to rapidly acclimate to different light conditions. The adaptive value of emigrating from the shade may thus deteriorate through time because the probability of acclimating to a different light intensity increases with the length of exposure (cells can begin acclimating to different light intensities within minutes, Bonente et al. 2012). The ‘doubling’ of costs associated with re-acclimating to light should reduce the penchant for movement by cells that have previously acclimated to shade.

The acclimation-cost hypothesis, which has similarities to Stamps’ ‘silver spoon’ and ‘natal habitat preference induction’ models (Davis & Stamps 2004; Stamps 2006; Stamps, Luttbeg & Krishnan 2009), likely accounts for why my best evidence for an ideal free distribution was associated with control dishes in which I observed no difference between halves in either density or fitness. Cells moving between identical ‘habitats’ do not change photosystems and glide through the media with normal flagellar movements that are unlikely to entail significant additional costs of habitat selection. Acclimation costs of habitat selection also likely account for the departure of the ‘released in light’ isodar from that predicted by data from control dishes. Cells migrating to shade must pay the cost of changing photosystems, but cells grown in light controls do not. This important caveat, that controls may not fully account for fitness expectations between habitats, should be carefully contemplated in future tests of habitat selection.

Regardless as to mechanisms, selection of high fitness habitat by *C. reinhardtii* in this study was conditional on whether nutrient concentrations were high (unused media) or low (used media). Although conditional strategies of habitat selection are inferior to density-dependent habitat choice, they can be adaptive when habitats of different quality remain constant over long periods of time (Morris, Diffendorfer, & Lundberg 2004). It will be interesting to learn whether

habitat selection by more sentient organisms also depends on differences between habitats conditioned by the mean quality of the environment.

Adaptive movement emphasizes the benefits of increased fitness as individuals select habitat in response to differences in their environment (Abrams 2000; Cressman & Křivan 2012). The distribution and fitness of *C. reinhardtii* in my experiments documents abilities of adaptive habitat choice originally developed mainly for sentient organisms (Fretwell & Lucas 1969; Flaxman & deRoos 2007). My research demonstrates rather clearly that even so-called simple single-celled organisms are capable of adaptive movements that modify spatial distribution and population dynamics. The biggest surprise, however, is not that algae are capable of adaptive habitat selection, but rather that adaptive movement is so often neglected by ecologists studying the dynamics of populations and communities.

## References

570 Abrams, P. A. 2000 The impact of habitat selection on the spatial heterogeneity of resources in  
571 varying environments. *Ecology* **81**, 2902–2913.

572 Bell, G. 1990 The ecology and genetics of fitness in *Chlamydomonas*. I. Genotype-by-  
573 environment interaction among pure strains. *Proceedings of the Royal Society of*  
574 *London B: Biological Sciences* **240**, 295–321.

575 Bonente, G., Pippa, S., Castellano, S., Bassi, R., & Ballottari, M. 2012 Acclimation of  
576 *Chlamydomonas reinhardtii* to different growth irradiances. *Journal of Biological*  
577 *Chemistry* **287**, 5833–5847.

578 Bold, H. C. 1949 The morphology of *Chlamydomonas chlamydogama* sp. nov. *Bulletin of the*  
579 *Torrey Botanical Club* **76**, 101–108.

580 Cressman, R. & Krivan, V. 2010 The ideal free distribution as an evolutionary stable state in  
581 density-dependent population games. *Oikos* **119**, 1231–1242.

582 Cressman, R. & Krivan, V. 2012 Two-patch population models with adaptive dispersal: the  
583 effects of varying dispersal speeds. *Journal of Mathematical Biology* DOI 10.1007/s00285-  
584 012-0548-3

585 Davis, J. M. & Stamps, J. A. 2004 The effect of natal experience on habitat preferences. *Trends*  
586 *in Ecology and Evolution* **19**, 411–416.

587 Ermilova, E., V., Zalutskaya, Z., M., Nikitin, M., M., Lapina, T. V., Fernández, E. 2010 R  
588 Regulation by light of ammonium transport systems in *Chlamydomonas reinhardtii*. *P*  
589 *Plant, Cell and Environment* **33**, 1049–1056.

- 590 Flaxman, S. M., & deRoos, C. A. 2007 Different modes of resource variation provide a critical  
591 test of ideal free distribution models. *Behavioral Ecology and Sociobiology* **61**, 877–886.
- 592 Fretwell, S. D. & Lucas, H. L. Jr. 1969 On territorial behaviour and other factors influencing  
593 habitat distribution in birds. I. Theoretical development. – *Acta Biotheoretica* **19**, 16–36.
- 594 Harris, E. H., Stern D. B., Witman, G. B. 2009 *The Chlamydomonas Sourcebook*. (ed. E. H.  
595 Harris) 2<sup>nd</sup> ed. Elsevier Inc. Oxford University Press, USA.
- 596 Holt, R. D. 1985 Population dynamics in two-patch environments: Some anomalous  
597 consequences of an optimal habitat distribution. *Theoretical Population Biology* **28**, 181–  
598 208.
- 599 Holt, R. D., Barfield, M. 2001 On the relationship between ideal free distribution and the  
600 evolution of dispersal. *Dispersal*. (ed. Jean Clobert) , pp. 83-96. Oxford University Press,  
601 USA.
- 602 Hugie, D. M., & Grand, T. C. 1998 Movement between patches, unequal competitors and the  
603 ideal free distribution. *Evolutionary Ecology* **12**, 1–19.
- 604 Johnson, M. L., & Gaines, M. S. 1990 Evolution of dispersal: Theoretical models and empirical  
605 tests using birds and mammals. *Annual Review of Ecology and Systematics* **21**, 449–480.
- 606 Křivan, V., Cressman, R., Schneider, C. 2008 The ideal free distribution: a review and synthesis  
607 of the game-theoretic perspective. *Theoretical Population Biology* **73**, 403–425.
- 608 Kuefler, D., Avgar, T., & Fryxell, J. M. 2013 Density- and resource-dependent movement  
609 characteristics in a rotifer. *Functional Ecology* **27**, 323–328.

- 610 Leboffe, M.J. & Pierce, B. E. 2005 *A photographic atlas for the microbiology laboratory*.  
 611 3<sup>rd</sup> ed. Morton Publishing Company, Colorado.
- 612 Milinski, M. & Parker, G. A. 1991 Competition for resources. In *Behavioural Ecology* (eds. J.  
 613 R. Krebs and N. B. Davies), 3<sup>rd</sup> edition. Blackwell Scientific, Oxford, UK.
- 614 Morris, D. W. 1987 Tests of density-dependent habitat selection in a patchy environment.  
 615 *Ecological Monographs* **57**, 269–281.
- 616 Morris, D. W. 1988 Habitat-dependent population regulation and community structure.  
 617 *Evolutionary Ecology* **2**, 253–269.
- 618 Morris, D. W. 2011 Adaptation and habitat selection in the eco-evolutionary process.  
 619 *Proceedings of the Royal Society of London B* **278**, 2401–2411.
- 620 Morris, D. W., Diffendorfer, J. E., Lundberg, P. 2004 Dispersal among habitats varying in  
 621 fitness: reciprocating migration through ideal habitat selection. *Oikos* **107**, 559–575.
- 622 Morris, D. W., Clark, R. & Boyce, M. S., 2008 Habitat and habitat selection: theory, test, and  
 623 implications. *Israel Journal of Ecology and Evolution* **54**, 287–294.
- 624 Poulin, M., Tuval, I., Drescher, K., Gollub, J. P., Goldstein, R. E. 2009 *Chlamydomonas* swims  
 625 with two “gears” in a eukaryotic version of run-and-tumble locomotion. *Science* **325**,  
 626 487–490.
- 627 R Development Core Team. 2008 *R: A language and environment for statistical computing*. R  
 628 Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, Available  
 629 online from <http://www.r-project.org/>.
- 630 Rosenzweig, M. L. 1981 A theory of habitat selection. *Ecology* **62**, 327–335.

631 Stamps, J. A. 2006 The silver spoon effect and habitat selection by natal dispersers. *Ecology*  
632 *Letters* **9**, 1179–1185

633 Stamps, J. A. 2009 Habitat selection. In *The Princeton Guide to Ecology*. (ed. S.A. Levin), pp.  
634 38–44. Princeton University Press, USA.

635 Stamps, J. A., Luttbeg, B. & Krishnan, V. V. 2009 Effects of survival on the attractiveness of  
636 cues to natal dispersers. *American Naturalist* **173**, 41–46.

637 Syrový, L., & Hodný, Z. 1991 Staining and quantification of proteins separated by  
638 polyacrylamide gel electrophoresis. *Journal of Chromatography* **569**, 175–196.

639 Tilman, D., 1990 Constraints and tradeoffs: Toward a predictive theory of competition and  
640 succession. *Oikos* **58**, 3–15.

641

642

643

644

645

646

647

648

649



## Appendix 1: DENSITY-DEPENDENT HABITAT SELECTION

Habitat selection refers to the process by which an individual chooses an area in which to conduct specific activities and interact with others (Stamps 2009). Habitat selection emerges because organisms are better suited to live and reproduce in some places than others (Morris et al. 2008). Individuals maximizing fitness should occupy the best habitat available. When increasing density depresses fitness to that of lower-quality habitats, individuals should disperse to those habitats ((Fretwell & Lucas 1969; Rosenzweig 1981; Morris 1987; Johnson & Gaines 1990; Holt & Barfield 2001). Thus, if individuals possess ‘complete’ knowledge of all habitat qualities, are of equal competitive ability, and if there is no cost to movement, then the distribution of individuals among habitats should fit an ideal free distribution (IFD) such that mean fitness is equal in every occupied habitat. If we assume logistic population growth such that

$$\frac{dN_i}{dt} = r_i N_i \left[ 1 - \frac{N_i}{K_i} \right] \quad (\text{A1})$$

where  $N_i$  is the population density,  $r_i$  is the maximum growth rate, and  $K_i$  is the carrying capacity in habitat  $i$ , and estimate fitness ( $W_i$ ) as per capita growth rate, then

$$W_i = \frac{1}{N_i} \frac{dN_i}{dt} = r_i - \frac{r_i N_i}{K_i} \quad (\text{A2})$$

(fitness declines linearly with increasing density, Fig. A1).

671

672

673

674

675

676

677

678

679 Figure A1: An illustration of ideal-free habitat selection with logistic population growth.

680 Ideal-free habitat selectors achieve equal expectations of fitness ( $W$ ) in habitats 1 and 2 (dashed  
681 horizontal lines), but at different population densities ( $N_1$  and  $N_2$ , dotted horizontal lines).

682

683

684

685

686

687

688

689


690

691

692

693

694



695

696

697

698

699

## Appendix 2: HABITAT ISODAR

Imagining that two habitats differ in logistic population growth, then the solution to ideal habitat selection is given by the habitat isodar (Morris 1987;1988)

$$N_2 = \frac{r_2 - r_1}{r_2} K_2 + \frac{r_1 K_2}{r_2 K_1} N_1 \quad (\text{A3})$$

(Fig. A2; Morris 1987; 1988). If one knows the rate at which fitness declines with density in two or more habitats, as in my experiments with *Chlamydomonas reinhardtii*, one simply needs to set the two fitness functions equal to one another in order to model the IFD isodar *a priori*. Knowing the isodar expected from ideal-free habitat selection enables a rigorous test for the ideal free distribution as long as one can then expose populations to the two habitats experimentally. The similarity between the predicted isodar and the distribution of individuals in the two-habitat experiments will reveal the ability of the organisms to achieve an ideal free distribution.

720

721

722

723

724

725

726

727

728 Figure A2: An example of an ideal-free habitat isodar (the distribution of individuals between  
729 two habitats such that mean fitness is equal in each) that emerges when fitness declines linearly  
730 with increasing density ( $N$ ) in habitats 1 and 2 as in figure A1. The dashed lines correspond to  
731 the carrying capacities in each habitat.

732

733

734


735

736

737

738

739  
740  
741  
742  
743



744  
745  
746  
747  
748

### **Appendix 3: BOLD'S BASAL MEDIUM**

I prepared modified Bold's basal medium for standard stock solution in four 1-litre batches (Table A1; Bold 1949; Bell 1990). I added the first six macronutrient salt solutions individually after complete dissolution into the medium, followed by trace metal and nutrient solutions (again, added individually after complete dissolution into the medium). I filtered ferrous sulphate heptahydrate through Whatman's filter paper No. 1, then autoclaved the filtered solution with the rest of the dissolved ingredients before combining them to create the complete medium. I adjusted the pH after autoclaving when necessary [pH = 6.7 (+/-0.2)].

772

773

774

775

776

777

778

779

780

781 Table A1: Stock solutions and volumes of each compound used in the recipe for modified

782 Bold's basal medium. Original stock solution and refinements are listed as in Bold (1949) and

783 Bell (1990).

784

785

786

787

788

789

790

791

792

793

794



795

796

Chemical compound	1 Litre Stock Solution	Quantity added to 1 litre of stock medium	Molar Concentration in final medium
<b>NaNO<sub>3</sub></b>	25 g L <sup>-1</sup> ddH <sub>2</sub> O	10 ml	2.94 x 10 <sup>-3</sup> M
<b>CaCl<sub>2</sub>·2H<sub>2</sub>O</b>	2.5 g L <sup>-1</sup> ddH <sub>2</sub> O	10 ml	1.7 x 10 <sup>-4</sup> M
<b>MgSO<sub>4</sub>·7H<sub>2</sub>O</b>	7.5 g L <sup>-1</sup> ddH <sub>2</sub> O	10 ml	3.04 x 10 <sup>-4</sup> M
<b>K<sub>2</sub>POH<sub>4</sub></b>	7.5 g L <sup>-1</sup> ddH <sub>2</sub> O	10 ml	4.31 x 10 <sup>-4</sup> M
<b>KH<sub>2</sub>PO<sub>4</sub></b>	17.5 g L <sup>-1</sup> ddH <sub>2</sub> O	10 ml	1.29 x 10 <sup>-3</sup> M
<b>NaCl</b>	2.5 g L <sup>-1</sup> ddH <sub>2</sub> O	10 ml	4.28 x 10 <sup>-4</sup> M
<b>EDTA anhydrous</b>	50 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	4.28 x 10 <sup>-4</sup> M
<b>KOH</b>	31 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	1.38 x 10 <sup>-3</sup> M
<b>FeSO<sub>4</sub>·7H<sub>2</sub>O</b>	4.98 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	4.48 x 10 <sup>-5</sup> M
<b>H<sub>2</sub>SO<sub>4</sub></b>	24.5 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	1 x 10 <sup>-3</sup> M
<b>H<sub>3</sub>BO<sub>3</sub></b>	11.42 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	4.62 x 10 <sup>-4</sup> M
<b>ZnSO<sub>4</sub>·7H<sub>2</sub>O</b>	8.82 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	7.67 x 10 <sup>-5</sup> M
<b>MnCl<sub>2</sub>·4H<sub>2</sub>O</b>	1.44 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	1.82 x 10 <sup>-5</sup> M
<b>MoO<sub>3</sub></b>	0.71 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	1.23 x 10 <sup>-5</sup> M
<b>CuSO<sub>4</sub>·5H<sub>2</sub>O</b>	1.57 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	1.57 x 10 <sup>-5</sup> M
<b>Co(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O</b>	0.49 g L <sup>-1</sup> ddH <sub>2</sub> O	1 ml	4.21 x 10 <sup>-6</sup> M

797

798

#### Appendix 4: QUANTIFYING CELL DENSITY


I calculated cell densities (millions of cells·ml<sup>-1</sup>) by fitting optical densities (spectrophotometer absorbancy) to haemocytometer cell counts by geometric mean regression. I used a dilution series to create different densities, then measured absorbancy at 665 *nm* in a microplate spectrophotometer (BioRad xMark™ Microplate Absorbance Spectrophotometer, 168-1150, CA, USA). I immobilized cells from corresponding samples in the two chambers of a Neubauer haemocytometer (Hausser Scientific, Catalogue # 3110, PA, USA) with a drop of Iodine-Potassium-Iodide (Lugol's stain, 6% KI, and 4% I).

I photographed the two chambers in the prepared haemocytometer slides separately with a microscope camera (OptixCam VS1.009, VA, USA), displayed the digital images on a computer monitor (Figure A7), then counted the number of cells in the four corner grids in each chamber's field-of-view. I used a total of 18 separate samples to calibrate optical densities for light and shade habitats at both the start and end (12 h) of the light cycle. Each estimate represents the mean number of cells counted from four separate images covering four separate sections of the haemocytometer grid. Methods for all of the haemocytometer cell counts were from Leboffe & Pierce (2005).

821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843

Figure A3: Regressions used to calibrate cell densities (millions of cells·ml<sup>-1</sup>) from optical densities (absorbancy at 665 nm) at the start of the light cycle for both light (a: open squares) and shaded (b: filled squares) habitats (*n* = 18 each). OD = optical density.

844  
845  
846  
847  
848  
849




850  
851  
852  
853  
854  
855  
856  
857

858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880

Figure A4: Regressions used to calibrate cell densities (millions of cells·ml<sup>-1</sup>) from optical densities (absorbancy at 665 nm) at the end of the light cycle (12 h) for both light (a: open squares) and shaded (b: filled squares) habitats (*n* = 18 each). OD = optical density.

881  
882  
883  
884  
885  
886  
887



888  
889  
890  
891  
892  
893  
894

895    **Appendix 5: LABORATORY & EXPERIMENTAL PROTOCOLS**

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940

Table A2: Culturing and experimental protocol used to assess habitat selection by  
*Chlamydomonas reinhardtii*. Experiments conducted between September 2012 and January  
2013.



Time	Culturing and experimental protocols
<b>Day 0</b>	Transfer single colony-forming units (CFUs) from enriched agar plates into 75 ml of Bold's medium in 250 ml Erlenmeyer flasks. Grow 'starting' cultures for 5-7 d to a mean density of 2.5 million cells·ml <sup>-1</sup> . Repeat for additional cultures as required. Grow all cultures in a controlled growth chamber at 23°C, set on a 12 h light:dark cycle to synchronize cellular division (Harris et al. 2009).
<b>Day 6-8</b>	Pipette 25-ml from each 'starting' culture into 300 ml of fresh medium in one litre Erlenmeyer flask for an additional growth period (7-10 d). Repeat for additional flasks.
<b>Day 14-18</b>	Adjust densities from batch cultures to three predetermined experimental densities by centrifugation and or dilution. Pelletize experimental cell cultures by centrifugation and re-suspend in 100 ml of fresh medium in 500 ml Erlenmeyer flasks for 36 h before use in order to acclimate cells to experimental conditions (Harris et al. 2009).
<b>Start of Experiment</b>	
<b>Time Zero (~day 20)</b>	Pipette 15 ml of experimental culture into one half of each partitioned Petri dish. Repeat with 4 dishes (one for each treatment) at three separate densities to be tested simultaneously. Pipette 15 ml of unused (or used media) medium into the alternate half of each Petri dish. Remove partitions from Petri dishes. Draw 8-220µl aqueous samples from each half of the dishes, parallel to the partition, and transfer into 96-well microplates. Record optical densities from spectrophotometer absorbencies at 665 nm. Overlay dishes with two layers of black nylon mesh for shade treatments. Place dishes in the incubator and do not disturb dishes for 12 h.
<b>Time 12</b>	Remove shade covers and insert new partitions along the central axis of each Petri dish and aerate each culture. Draw 8-220µl aqueous samples from each half of the dishes parallel to the partition, and transfer into 96-well microplates. Record optical densities from spectrophotometer absorbencies at 665 nm. Reposition covers for shade treatments.
<b>Time 24</b>	Repeat sampling. Remove shade cover, then aerate and sample cell densities in each half of each Petri dish. Draw 8-220µl aqueous samples from each half of the dishes perpendicularly to the partition, and transfer into 96-well microplates. Read optical densities on the spectrophotometer at 665 nm. Reposition shade cover and then repeat with next dish.
<b>Time 48</b>	Remove shade covers and insert new partitions along central axis of Petri dish and aerate culture in each half of the dishes. Draw 8-220µl aqueous samples from each half of the dishes perpendicular to the partition, and transfer into 96-well microplates. Record optical densities from spectrophotometer absorbencies at 665 nm.
<b>End of experiment</b>	

943    **Appendix 6: PHOTOGRAPHS**

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963


964

965

966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988

Figure A5: Photograph illustrating an example of the random placement of control (fully covered by shade and fully exposed to light) and habitat-selection Petri dishes (half shaded) used to assess habitat selection by *Chlamydomonas reinhardtii*. Each of the three replicate dishes contains a different algal density. Image taken at 12 h.

989  
990  
991  
992  
993  
994  
995



996  
997  
998  
999  
1000  
1001  
1002

1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025

Figure A6: Photograph of a Petri dish illustrating habitat selection by *Chlamydomonas reinhardtii* cells released in the light habitat (right-hand side) with opportunity to occupy shade (unused media). Initial density at time zero = 0.85 million cells·ml<sup>-1</sup>. Image taken at 12 h.

1026

1027


1028

1029

1030

1031

1032



1033

1034

1035

1036

1037

1038

1039

1040

1041

1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064

Figure A7: Photograph of immobilized *Chlamydomonas reinhardtii* cells in a haemocytometer chamber used to calibrate optical densities. Living cells are dark green. Dead cells (not counted) appear without complete cellular structure or vibrancy (Leboffe & Pierce 2005).


1065

1066

1067

1068

1069



1070

1071

1072

1073

1074

1075



1076    **Appendix 7: DIFFUSION**

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121

Table A3: Comparisons of paired mean absorbancy values between the initial (dye added) and alternate (no dye) sides of Petri dishes at 12 hourly intervals. Degrees of freedom at 3 and 11 hours were reduced because I replaced one randomly chosen sample with a pure-media blank in order to calibrate the spectrophotometer for each of the two 96-well microplates evaluating absorbancy of the 192 samples (16 samples  $\times$  12 hours). Bold lettering identifies statistically significant differences. Paired t-tests; two-tailed significance.

Hour	Side	Mean OD	Paired <i>T</i>	df	<i>P</i>
0	Initial	0.059	-9.23	7	<0.001
	Alternate	0.017			
1	Initial	0.048	9.45	7	<0.001
	Alternate	0.037			
2	Initial	0.056	7.86	7	<0.001
	Alternate	0.028			
3	Initial	0.052	9.65	6	<0.001
	Alternate	0.025			
4	Initial	0.044	2.34	7	0.054
	Alternate	0.040			
5	Initial	0.048	1.95	7	0.09
	Alternate	0.045			
6	Initial	0.044	-1.79	7	0.12
	Alternate	0.048			
7	Initial	0.046	-1.48	7	0.18
	Alternate	0.048			
8	Initial	0.044	0.04	7	0.97
	Alternate	0.044			
9	Initial	0.043	0.55	7	0.60
	Alternate	0.041			
10	Initial	0.043	0.10	7	0.928
	Alternate	0.043			
11	Initial	0.048	2.07	6	0.08
	Alternate	0.043			